

On radar time and the twin “paradox”

Carl E. Dolby^{a)} and Stephen F. Gull^{b)}

Astrophysics Group, Cavendish Laboratory, Madingley Road, Cambridge CB3 0HE, United Kingdom

(Received 23 April 2001; accepted 13 June 2001)

In this paper we apply the concept of radar time (popularized by Bondi in his work on k calculus) to the well-known relativistic twin “paradox.” Radar time is used to define hypersurfaces of simultaneity for a class of traveling twins, from the “immediate turn-around” case, through the “gradual turn-around” case, to the “uniformly accelerating” case. We show that this definition of simultaneity is independent of choice of coordinates, and assigns a *unique* time to any event (with which the traveling twin can send and receive signals), resolving some common misconceptions.

© 2001 American Association of Physics Teachers.

[DOI: 10.1119/1.1407254]

INTRODUCTION

It might seem reasonable to suppose that the twin paradox has long been understood, and that no confusion remains about it. Certainly there is no disagreement about the relative aging of the two twins, so there is no “paradox.” There is also no confusion over “when events are seen” by the two twins, or over the description by the stay-at-home twin (Alex, say), of “when events happened.” These aspects of the twin paradox are treated in the standard texts.^{1–4}

However, the description of “when events happened” according to the traveling twin (Barbara, say) seems never to have been fully settled.^{5,6} In textbook treatments, Barbara’s hypersurfaces of simultaneity, which define “when events happened” according to her, have consistently been misrepresented or ignored. A common diagram for such hypersurfaces^{1,2,3,6} is shown in Fig. 1.

Bohm¹ and Sartori³ claim that points G and H appear to the traveling twin as simultaneous, while Bohm¹ goes on to say that “after the acceleration at E an event such as N is ascribed a smaller time coordinate than it had before!” Other authors,^{3,2} noticing this problem, claim that its resolution lies in an analysis of the interval that the traveling twin must take to turn around (at a finite acceleration), and that during this period of acceleration Barbara’s hypersurfaces of simultaneity “sweep round” from EG to FH , as shown in Fig. 2.

Although this period of acceleration can indeed fix the gap between G and H , it cannot resolve the more serious problem (mentioned also in Marder⁷ and in Misner *et al.*⁸) which occurs to Barbara’s left. Here her hypersurfaces of simultaneity are overlapping, and she assigns three times to every event! Also, if Barbara’s hypersurfaces of simultaneity at a certain time depend so sensitively on her instantaneous velocity as these diagrams suggest, then she would be forced to conclude that the distant planets swept backwards and forwards in time whenever she went dancing!

The path through this confusion existed already in Einstein’s original paper,⁹ and was popularized by Bondi in his work on “ k calculus.” It lies in the correct application of “radar time” (referred to as “Märzke-Wheeler Coordinates” in Pauri *et al.*¹⁰). This concept is not new. Indeed, Bohm¹ and D’Inverno² both devote a whole chapter to k calculus, and use “radar time” (not under that name) to derive the hypersurfaces of simultaneity of inertial observers. However, both authors then apply this definition wrongly to the traveling twin.

In the present paper we recall the definition of “radar

time” (and related “radar distance”) and emphasize that this definition applies not just to inertial observers, but to any observer in any space–time. We then use radar time to derive the hypersurfaces of simultaneity for a class of traveling twins, from the “immediate turn-around” case, through the “gradual turn-around” case, to the “uniformly accelerating” case. (The “immediate turn-around” and “uniformly accelerating” cases are also discussed in Pauri *et al.*¹⁰) We show that in all cases this definition assigns a *unique* time to any event with which Barbara can send and receive signals, and that this assignment is independent of any choice of coordinates. We then demonstrate that brief periods of acceleration have negligible effect on the radar time assigned to distant events, in contrast to the sensitive dependence of the hypersurfaces implied by Figs. 1 and 2. By viewing the situation in different coordinates we further demonstrate the coordinate independence of radar time, and note that there is no observational difference between the interpretations in which the differential aging is “due to Barbara’s acceleration” or “due to the gravitational field that Barbara sees because of this acceleration.”

RADAR TIME AND RADAR DISTANCE

Consider an observer traveling on path $\gamma: x^\mu = x^\mu_{(\gamma)}(\tau)$ with proper time τ (Fig. 3). Define:

$\tau^+(x) \equiv$ (earliest possible) proper time at which
a light ray (technically, a null geodesic)
leaving point x could intercept γ ,
 $\tau^-(x) \equiv$ (latest possible) proper time at which a
light ray (null geodesic) could leave γ ,
and still reach point x ,

$$\tau(x) \equiv \frac{1}{2}(\tau^+(x) + \tau^-(x)) = \text{“radar time,”}$$

$$\rho(x) \equiv \frac{1}{2}(\tau^+(x) - \tau^-(x)) = \text{“radar distance,”}$$

$$\Sigma_{\tau_0} \equiv \{x: \tau(x) = \tau_0\}$$

= observer’s “hypersurface

of simultaneity at time τ_0 .”

This definition was made popular by Bondi in his work on special relativity and k calculus,^{11,12} and is essentially a re-




Fig. 1. A typical “textbook illustration” of the hypersurfaces of simultaneity of the traveling twin in the twin paradox.

arrangement of the formula $t - t^- = t^+ - t$ from Einstein’s original paper⁹ (written as $t_B - t_A = t'_A - t_B$ in Einstein’s notation). This formula simply says that an observer can assign a time to a distant event by sending a light signal to the event and back, and averaging the (proper) times of sending and receiving. It is clearly applicable to any observer in any gravitational background. However, the idea of defining hypersurfaces of simultaneity in terms of “radar time” has seldom been applied to noninertial observers or to nonflat space-times. This is perhaps due to Bondi’s claim that¹¹ “how a clock reacts to acceleration is utterly dependent on how the clock is constructed.” This claim has since been refuted by experiment (Ref. 7, or Ref. 12, and pp. 144 and 145, and references therein); moreover the assumption that “suitable clocks” will behave identically under acceleration




Fig. 2. Another typical “textbook illustration,” in which Barbara’s hypersurfaces of simultaneity “sweep round” from EG to FH during her period of acceleration.




Fig. 3. Schematic of the definition of “radar time” $\tau(x)$.

(or under gravitational fields) is a basic premise of general relativity, without which proper time would have no physical meaning. There is therefore no reason not to apply “radar time” to accelerating observers or to curved space-times.

Since radar time can be defined without any mention of coordinates it is, by construction, independent of our choice of coordinates. It is single valued, agrees with proper time on the observer’s path, and is invariant under “time reversal”—that is, under reversal of the sign of the observer’s proper time.

THE TRAVELING TWIN

Consider the standard “immediate turn-around” case, for which Barbara’s path is given by

$$(t_B(\tau), x_B(\tau)) = \begin{cases} (\cosh(\alpha)\tau, \sinh(\alpha)\tau) & (\tau > 0) \\ (\cosh(\alpha)\tau, -\sinh(\alpha)\tau) & (\tau < 0) \end{cases}$$

or

$$(u_B(\tau), v_B(\tau)) = \begin{cases} (e^\alpha \tau, -e^{-\alpha} \tau) & (\tau > 0) \\ (e^{-\alpha} \tau, -e^\alpha \tau) & (\tau < 0) \end{cases}$$

where $s = \tanh(\alpha)$ is Barbara’s speed (in units with $c = 1$), α is her “rapidity,” and we have defined coordinates (u, v) by $u \equiv x + t$ and $v \equiv x - t$. This situation is shown in Fig. 4.




Fig. 4. The traveling twin in the “immediate turn-around” case.




Fig. 5. Hypersurfaces of simultaneity of the traveling twin in the “immediate turn-around” case.

To deduce the radar time of a point (u_0, v_0) , it is convenient to consider four different regions, depending on the signs of u_0 and v_0 .

Region I: $u_0 > 0$ and $v_0 > 0$. Clearly, $\tau^+(u_0, v_0)$ is determined by putting $u_0 = u_B(\tau^+) = e^\alpha \tau^+$ [where we have used $u_B(\tau^+) = e^\alpha \tau^+$ since we see from Fig. 4 that $\tau^+ > 0$]. Similarly, $\tau^-(u_0, v_0)$ is determined by putting $v_0 = v_B(\tau^-) = -e^\alpha \tau^-$ (since $\tau^- < 0$). Hence:

$$\tau(u_0, v_0) = \frac{1}{2}(e^{-\alpha} u_0 - e^{-\alpha} v_0) = e^{-\alpha} t_0 \quad (1)$$

in this region.

Region F: $u_0 > 0$ and $v_0 < 0$. Consider first the case where (u_0, v_0) is to Barbara’s right, as shown in Fig. 4. The calculation of $\tau(u_0, v_0)$ proceeds as in the previous case, the only difference being that, since $\tau_- > 0$, we now use $v_B(\tau_-) = -e^{-\alpha} \tau_-$ (the exponent has changed sign). Hence we have

$$\tau(u_0, v_0) = \frac{1}{2}(e^{-\alpha} u_0 - e^\alpha v_0) = \cosh(\alpha) t_0 + \sinh(\alpha) x_0. \quad (2)$$

If (u_0, v_0) is to Barbara’s left (but still in region F) then the roles played by τ^+ and τ^- are reversed, but since the definition of τ is symmetric under this change it follows that (2) remains unchanged, and holds throughout region F.

Regions II and P are treated similarly, and we find

$$\tau(t, x) = \begin{cases} t e^{-\alpha}, & \text{region I} \\ t e^\alpha, & \text{region II} \\ \cosh(\alpha) t + \sinh(\alpha) x, & \text{region F} \\ \cosh(\alpha) t - \sinh(\alpha) x, & \text{region P} \end{cases}$$

The “hypersurface of simultaneity at time τ_0 ,” denoted $t_{\tau_0}(x)$, is now given by putting $\tau(t, x) = \tau_0$ and rearranging. This gives

$$t_{\tau_0}(z) = \begin{cases} \tau_0 e^\alpha, & \text{region I} \\ \tau_0 e^{-\alpha}, & \text{region II} \\ \text{sech}(\alpha) \tau_0 - \tanh(\alpha) x, & \text{region F} \\ \text{sech}(\alpha) \tau_0 + \tanh(\alpha) x, & \text{region P} \end{cases}$$

as shown for various τ_0 in Fig. 5.

We now see that in region F, where Barbara is indistinguishable from an inertial observer with velocity s , her hy-

persurfaces of simultaneity are exactly those appropriate to such an inertial observer; while in region P, where she is indistinguishable from an inertial observer with velocity $-s$, her hypersurfaces are those appropriate to that inertial observer. This is not true in regions I and II however, for which she has differing velocities when she sends her signal (at time τ^-) and when she receives it back (at time τ^+). In these regions her hypersurfaces of simultaneity are flat (as expected for symmetry under time reversal). Hence she assigns a *unique* time to each event, avoiding the problems that appear in the common “textbook illustration (Figs. 1 and 2).”

We now consider the effect on these conclusions of a small period of acceleration at the point of turn around.

GRADUAL TURN-AROUND

We allow Barbara to turn around at finite acceleration a , for a proper time $\tau_c = \alpha/a$. Barbara’s trajectory is now given by

$$t_B(\tau) = \begin{cases} \cosh(a\tau_c)(\tau - \tau_c) + \frac{1}{a} \sinh(a\tau_c) & (\tau \geq \tau_c) \\ \frac{1}{a} \sinh(a\tau) & (|\tau| \leq \tau_c) \\ \cosh(a\tau_c)(\tau + \tau_c) - \frac{1}{a} \sinh(a\tau_c) & (\tau \leq -\tau_c) \end{cases},$$

$$x_B(\tau) = \begin{cases} \sinh(a\tau_c)(\tau - \tau_c) + \frac{1}{a} \cosh(a\tau_c) & (\tau \geq \tau_c) \\ \frac{1}{a} \cosh(a\tau) & (|\tau| \leq \tau_c) \\ -\sinh(a\tau_c)(\tau + \tau_c) + \frac{1}{a} \cosh(a\tau_c) & (\tau \leq -\tau_c) \end{cases},$$

and her final speed is $s = \tanh(a\tau_c)$. In null coordinates this is

$$(u_B(\tau), v_B(\tau)) = \begin{cases} \left(e^{a\tau_c} \left(\tau - \tau_c + \frac{1}{a} \right), -e^{-a\tau_c} \left(\tau - \tau_c - \frac{1}{a} \right) \right) & (\tau \geq \tau_c) \\ \left(\frac{1}{a} e^{a\tau}, \frac{1}{a} e^{-a\tau} \right) & (|\tau| \leq \tau_c) \\ \left(e^{-a\tau_c} \left(\tau + \tau_c + \frac{1}{a} \right), -e^{a\tau_c} \left(\tau + \tau_c - \frac{1}{a} \right) \right) & (\tau \leq -\tau_c) \end{cases}.$$

We now have nine regions to consider, separated by $u = e^{\pm a\tau_c}$ and $v = e^{\pm a\tau_c}$. The calculation is straightforward, and yields hypersurfaces of simultaneity of the form:

$$t_{\tau_0}(x) = \begin{cases} \tau_0 e^{\pm a\tau_c}, & \text{regions I, II} \\ \text{sech}(a\tau_c)(\tau_0 \pm \tau_c) \mp \tanh(a\tau_c)x, & \text{regions P, F} \\ x \tanh(a\tau_0), & \text{region U} \end{cases}$$

while in the other four regions these hypersurfaces are given implicitly by

$$\log(a|x + t_{\tau_0}(x)|) + a(t_{\tau_0}(x) - x)e^{\pm a\tau_c} = 2a\tau_0 \mp a\tau_c - 1, \quad \text{regions FII, PI}$$




Fig. 6. Hypersurfaces of simultaneity of the traveling twin in the “gradual turn-around” case.

$$\log(a|x - t_{\tau_0}(x)|) - a(t_{\tau_0}(x) + x)e^{\pm a\tau_c} = -2a\tau_0 \mp a\tau_c - 1, \text{ regions FI, PII.}$$

These hypersurfaces are shown in Fig. 6. As hoped, there is very little difference (for small τ_c) between the “immediate turn-around” and “gradual turn-around” cases. The hypersurfaces in regions I and II are identical to the previous case, while in regions F and P the only difference is that a slightly different value of τ_0 is assigned to each hypersurface, due to the slight difference in turn-around time. This contrasts with the situation described in many textbooks^{7,2} and depicted in Fig. 2.

It is interesting to consider the limit $\tau_c \rightarrow \infty$, corresponding to a uniformly accelerating observer. In this case we have simply

$$(t_B(\tau), x_B(\tau)) = \left(\frac{1}{a} \sinh(a\tau), \frac{1}{a} \cosh(a\tau) \right),$$

or

$$(u_B(\tau), v_B(\tau)) = \left(\frac{1}{a} e^{a\tau}, \frac{1}{a} e^{-a\tau} \right).$$

A simple calculation yields radar time and radar distance:

$$\tau(t, x) = \frac{1}{2a} \log\left(\frac{x+t}{x-t}\right),$$

$$\rho(t, x) = \frac{1}{2a} \log(a^2(x^2 - t^2)).$$

The hypersurfaces of simultaneity are therefore given by

$$t_{\tau_0}(x) = x \tanh(a\tau_0)$$

just as in region U above.

ρ and τ are simply Rindler coordinates now, and cover only region U. This is as expected, since Barbara cannot send a signal to any events in regions P or II, while no signals could reach Barbara from regions F or II.




Fig. 7. Hypersurfaces of simultaneity of a uniformly accelerating observer.

GRAVITY DOESN'T MATTER

It is often said of the twin paradox that⁴ “a complete explanation of the problem can only be given within the framework of general relativity.” However, as we have just shown, Barbara’s hypersurfaces of simultaneity depend only on the kinematics involved, and can be fully understood without resorting to general relativity. To understand why “equivalent gravitational fields” are sometimes invoked in the literature, consider the “uniformly accelerating” case, as described in (τ, ρ) coordinates. This is shown in Fig. 8.

The metric in these coordinates is $ds^2 = e^{2a\rho}(d\tau^2 - d\rho^2)$. Although Barbara has a “straight trajectory” in these coordinates, she is still undergoing an acceleration a , just as we undergo a constant acceleration g through being “held up” against the earth’s gravitational field. (Of course Barbara is here being “pulled down” rather than pulled “toward the earth’s center,” so no tidal forces are involved.) Although Alex’s trajectory appears curved in these coordinates, he is still inertial, since his trajectory is geodesic in this metric. Alex and Barbara’s proper times are unaltered by this change of coordinates, so that it is still Alex who ages more during his absence from Barbara. Some of the lines of constant t , which represent Alex’s hypersurfaces of simultaneity, fail to intercept his trajectory in Fig. 8. This is because Barbara’s (τ, ρ) coordinates cover only region U of Fig. 7, so any line of constant t which intercepts Alex’s trajectory in region F or P will not have the interception visible in Fig. 8.




Fig. 8. Alex’s hypersurfaces of simultaneity as described in Barbara’s (τ, ρ) coordinates, for the “uniformly accelerating” case.




Fig. 9. Alex's hypersurfaces of simultaneity, as described in Barbara's (τ, ρ) coordinates, for the "immediate turn-around" case.

A similar coordinate transformation can be performed for the "immediate turn-around" case. In (τ, ρ) coordinates this is shown in Fig. 9.

The metric in these coordinates is

$$ds^2 = \begin{cases} e^{-2\alpha}(d\tau^2 - d\rho^2), & \text{region P} \\ (d\tau^2 - d\rho^2), & \text{regions I, II} \\ e^{2\alpha}(d\tau^2 - d\rho^2), & \text{region R} \end{cases}$$

which is Minkowski in all four regions, but with a shock-like scale discontinuity along the lines $x = \pm t$ which causes geodesics to "turn left" upon crossing these lines. In these coordinates Alex's hypersurfaces of simultaneity look much like Barbara's did in the original coordinate system. It is still Alex who is inertial, however. Graphing the results in different coordinate systems makes no difference to this conclusion. Also, since the definition of radar time is independent of the choice of coordinates, then any statement that Barbara




Fig. 10. Hypersurfaces of simultaneity of the traveling twin in the more realistic case, in which their trajectories coincide before and after the journey.

makes of the form "Alex did this at this time" (according to her), or any statement that Alex makes of the form "Barbara did this at this time" (according to him), will be unchanged by changes in coordinates.

CONCLUSION

We have demonstrated that "radar time" is applicable to arbitrarily moving observers, and we have used it to find the hypersurfaces of simultaneity of the traveling twin in the "twin paradox." Radar time allows the traveling twin to assign times to distant events in a fully coordinate invariant fashion. We have considered a general class of twins, varying from the "immediate turn-around" case to the "uniformly accelerating" observer, and have highlighted some common misconceptions.

Throughout this paper we have made a convenient but unrealistic simplification. It is clear from the trajectories shown in the previous figures that Barbara and Alex could not possibly be twins, since they never spend more than an instant in each other's company. More realistically, Barbara's trajectory should coincide with Alex's until she begins her journey, and should coincide again after her return. The hypersurfaces of simultaneity for this more realistic trajectory are shown in Fig. 10.

In the *causal envelope*¹⁰ of Barbara's journey (that is: the intersection of the causal future of her departure and the causal past of her return, being the middle four regions in Fig. 10) these hypersurfaces are identical to those of Fig. 5. Readers might enjoy interpreting the rest of this diagram, convincing themselves that the length of the left arrow represents Barbara's traveling time and that the length of the middle arrow represents Alex's, and verifying that the ratio of these is indeed $\sqrt{1-s^2}$.

ACKNOWLEDGMENTS

We thank Dr. Anton Garrett for his many helpful comments on this manuscript. CD thanks Dr. Henry Levy for his enthusiastic teaching of k calculus, and thanks Cavendish Astrophysics, and Trinity College, Cambridge, for financial support.

^aElectronic mail: c.dolby@mrao.cam.ac.uk

^bElectronic mail: steve@mrao.cam.ac.uk

¹D. Bohm, *The Special Theory of Relativity* (W. A. Benjamin, New York, 1965).

²R. D'Inverno, *Introducing Einstein's Relativity* (Oxford U.P., Oxford, 1992).

³L. Sartori, *Understanding Relativity: A Simplified Approach to Einstein's Theories* (University of California Press, Berkeley, 1996).

⁴W. Pauli, *Theory of Relativity* (Pergamon, New York, 1958).

⁵W. G. Unruh, "Parallax distance, time, and the twin 'paradox,'" *Am. J. Phys.* **49** (6), 589–592 (1981).

⁶T. A. Debs and M. L. G. Redhead, "The twin 'paradox' and the conventionality of simultaneity," *Am. J. Phys.* **64** (4), 384–392 (1995).

⁷L. Marder, *Time and the Space-Traveller* (George Allen & Unwin Ltd., London, 1971).

⁸C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).

⁹A. Einstein, "On the Electrodynamics of Moving Bodies," *Ann. Phys. (Leipzig)* **17**, 891–921 (1905).

¹⁰M. Pauri and M. Vallisneri, "Märzke–Wheeler Coordinates for Accelerated Observers in Special Relativity," *Found. Phys. Lett.* **13** (5), 401–425 (2000).

¹¹H. Bondi, *Assumption and Myth in Physical Theory* (Cambridge U.P., Cambridge, 1967).

¹²P. C. W. Davies, *About Time* (Orion, London, 1995).